## **GPAT QUESTION PAPER 2020 WITH ANSWER KEY**

1. Relative sweetness of sucrose, to saccharin:

(a) 121°C for 15 mins (b) 121°C for 30 mins

(a) 1: 200	(b) 1: 500	(c) 1: 100	(d) 1: 400			
2. Following s prediction :	tatement is mo	re accurate wit	th respect to limitations of Arrhenius relationship for stability			
(a) Order of de	gradation will	alter at higher	temperature			
(b) Equal mois	sture concentra	tions will be m	nentioned at different temperatures			
(c) Less relativ	e humidity an	d oxygen solub	pility at higher temperature			
(d) Same degra	adation mecha	nisms may pre	dominate at different temperatures			
3. Lakes of Dy	es available co	ommercially co	ontain maximum upto of pure dye.			
(a) 15%	(b) 10%	(c) 25%	(d) 50%			
4. Type II glas	s containers ar	re				
(a) Suitable for	r alkaline solut	tions				
(b) Most inert	glasses and sh	ows high hydro	olytic resistance			
(c) Suitable for	r most acidic a	nd neutral aqu	eous preparations			
(d) Suitable fo	r non-aqueous	preparations				
•		•	nhancer for topical formulations by			
(a) Increasing			enaturing proteins			
(c) Increasing	transepiderma	l loss (d) Al	tering solvent nature of membrane			
6. As per Euro called very hys			al guide, substance stored at 25°C for 24 hours at 80% RH, reight is			
(a) 0.2% w/w a	(a) $0.2\%$ w/w and $<15\%$ w/w (b) $>0.2\%$ w/w and $<20\%$ w/w					
(c) > 15%  w/w	<b>V</b>	(d) 0.2	2% w/w and < 2% w/w			
7. SCHICK Te	est Toxin is a s	terile Filtrate f	rom a culture of			
(a) Rickettsia	prowazeki	(b) N	Mycobacterium Diphtheriae			
(c) Coryne bac	terium Diphth	eriae (d) A	Actinobacillus mallei			
8. Violin gut is	s obtained fron	n intestine of				
(a) Horse (b) C	Cat (c) Sheep (	d) Camel				
9. An excipien	t Ludipress us	ed for liquid D	osage form is a coprocessed excipient of			
(a) PticInsi. De	extrose monoh	ydrate + Kollid	don 30			
(b) Lactose mo	onohydrate + K	Collidon 30				
(c) Lactose monohydrate + Kollidon 30 + Kollidon CL						
(d) Dextrose m	nonohydrate +	Kollidon 30 +	Kollidon CL			
10. Pyrogens p	oresent in conta	ainers can be d	estroyed by heating the containers at			

(c) 210°C for 4 h
(d) 210°C for 1 h
11. In capsule making the Bloom strength of gelatin is proportional to molecular weight of the gelatin and is a measure of the;
(a) Cohesive strength of the solvent molecules
(b) Cohesive strength of the crosslinking that occurs between gelatin molecules
(c) Adhesive strength of gelatin with dipping pins
(d) Adhesive strength of gelatin with other polymer
12. Choose the wrong statement from the following with regard to Amorphous solids
(a) Usually they are anisotropic
(b) They tend to flow when subjected to sufficient pressure
(c) Considered as super cooled fluids
(d) They do not have definite melting point
13. The 'Crenulation' effect on the cell wall is caused by solutions and this effect is
(a) Hypertonic; reversible
(b) Hypotonic; reversible
(c) Hypotonic; irreversible
(d) Hypertonic; irreversible
14. The unequal attractive forces acting on the molecules at the surface of liquid gas interface when compared with molecular forces in the bulk of the liquid is due to:
(a) Abscence of adhesive force of attraction
(b) Less adhesive force of attraction
(c) Abscence of cohesive force of attraction
(d) Less cohesive force of attraction
15. Following is the important sterol in faeces formed from cholesterol by bacteria in lower intestine:
(a) 7-α-Hydroxy cholesterol
(b) Coprostanol
(c) 7-Dehydrocholesterol
(d) Lithocholic acid
16. According to distribution law, select the appropriate expression for the concentration of a solute when it exist as monomer in solvent A and dimer in solvent B (Assume CA = concentration of solute in solvent A CB = concentration of solute in solvent B $\alpha$ = degree of dissociation)
(a) $K = \frac{C_A}{2 \times \sqrt{C_B}}$ (b) $K = \frac{C_A}{C_B \times (1 - \alpha)}$ (c) $K = \frac{C_A}{C_B \times (2 - \alpha)}$ (d) $K = \frac{C_A}{\sqrt{C_B}}$

18. Production of Acetyl methyl carbinol can be detected by which of the following test

(b) Therapeutic index

concentration (MSC) is called


(a) Toxic range

17. The drug concentration between Minimum effective concentration (MEC) and maximum safe

(c) Therapeutic ratio (d) Therapeutic range

(a) Voges-proskaver test (b) Ind	dole test (c)	Citrate utilization	test (d) Methyl red test		
19. The temperature dependence and	the theory of visco	sity is expressed b	by the following		
(a) Stern-Volmer equation	(b) Arrhen	ius equation of che	emical kinetics		
(c) Higuchi equation	(d) Gibb's	equation			
20. Which of the following is not a f the construction of pharmaceutical o		y) factor consider	ed for selection of a location for		
(a) Soil (b) Market for produc	ets (c) Labor	supply (d) Ra	aw materials		
		tive for the coagul	ase test (a test for the ability of		
(a) S. epidermidis (b) S. aureus	(c) S. Saprophytic	us (d) S. lactis			
22. The version of GMP in India that cosmetics	t describe requirem	ent of factory pren	nises for manufacture of		
(a) schedule-M (b) Schedule-	M-III (c) Schedu	le-M-II (d) So	hedule M-I		
23. This fungus is also known as opp	oortunistic pathoger	, often isolated fro	om warm-blooded animals.		
(a) Aspergillus niger (b) Penicillinu	ım notatum (c)	Penicillinium chry	/sogenum (d) Candida albican		
24. Which of the following will resu	lt in very closest va	lue to the Glomeri	ılar Filteration Rate (GFR)?		
(a) Insulin Clearance	(b)	Albumin Clearand	e		
			operty of drug include		
26. The Acidic polymers of Ribitol/G known as	Glycerol phosphate	present in Gram p	ositive microorganism are		
(a) Polysaccharides (b) Teichoic a	acids (c) Peptide	oglycans (d) Ly	sozymes		
•			e than 60% may lead to		
	with the first transfer and pro-				
(a) $\Delta S$ universe = $S$ system + $\Delta S$ sur	rounding				
(b) S system = $\Delta$ S universe + $\Delta$ S sur	rounding				
(c) $\Delta S$ universe = $\Delta S$ surrounding - $S$	S system				
(d) $\Delta S$ universe = S system - $\Delta S$ sur	rounding				
29. The complexation of Quinine wi	th hexyl salicylate i	s an example of al	osorption by		
a) Soil (b) Market for products (c) Labor supply (d) Raw materials  1. The following gram positive bacteria species is positive for the coagulase test (a test for the ability of acteria to cause blood plasma to clot) a) S. epidermidis (b) S. aureus (c) S. Saprophyticus (d) S. lactis  2. The version of GMP in India that describe requirement of factory premises for manufacture of osmetics a) schedule-M (b) Schedule-M-III (c) Schedule-M-II (d) Schedule M-I  3. This fungus is also known as opportunistic pathogen, often isolated from warm-blooded animals. a) Aspergillus niger (b) Penicillinium notatum (c) Penicillinium chrysogenum (d) Candida albicans 4. Which of the following will result in very closest value to the Glomerular Filteration Rate (GFR)? a) Insulin Clearance (b) Albumin Clearance b) Measure of Blood Urea Nitrogen (BUN) (d) Creatinine Clearance b) Measure of Blood Urea Nitrogen (BUN) (d) Creatinine Clearance b) Heat capacity (b) Solubility (c) Refractive index (d) Entropy b) 6. The Acidic polymers of Ribitol/Glycerol phosphate present in Gram positive microorganism are nown as b) Polysaccharides (b) Teichoic acids (c) Peptidoglycans (d) Lysozymes 7. Increasing the proportion of the disperse phase of an emulsion by more than 60% may lead to a) Cracking (b) Frothing (c) Phase inversion (d) Creaming b) Cracking (b) Frothing (c) Phase inversion (d) Creaming c) As universe = S system + ΔS surrounding c) S system = ΔS universe + ΔS surrounding c) S system = ΔS universe + ΔS surrounding c) S system = ΔS universe + ΔS surrounding c) S system = ΔS universe + ΔS surrounding c) The complexation of Quinine with hexyl salicylate is an example of absorption by a) Convective transport (b) Facilitated transport (c) Penocytosis (d) Ion pair transport 0. The polymorphs exhibit the following different properties Except: a) X-ray crystal and diffraction patterns (b) Melting points c) O The polymorphs exhibit the following different properties Except: a) X-ray crystal and diffraction patterns (b) Melting points c) O The po					
30. The polymorphs exhibit the follo	wing different prop	erties Except:			
(a) X-ray crystal and diffraction patt	erns (b) Melting	g points			
(c) Solubilities	(d) Chemic	al structures			
31. Method of Study Drug Distribution pattern is :					
(a) Mass spectrophotometry	(b) Nuclea	r magnetic resonar	nce		
(c) Flourimetry	(d) UV-vis	ible spectrophotor	netry		

## 32 Product of the following



reaction:

- 33. Which of the following powder shows the greatest degree of dustibility
- (a) Talcum powder (b) Lycopodium powder (c) Fine charcoal (d) Potato Starch
- 34. The dispersion of course material by shearing in a narrow gap between a static cone and a rapidly rotating cone is caused by
- (a) Colloid Milk
- (b) Electrical Dispersion
- (c) Peptisation
- (d) Ultrasonic Irradiation
- 35. When the concentration of an aqueous sodium chloride solution has the same colligative properties as the solution in question, the value so obtained is known as
- (a) Normality (b) Isotonicity value (c) Molarity (d) Molality
- 36. When adhesive attractions between molecules of different species exceeds cohesive attraction between like molecules, the deviation according to Raoult's law is said to be
- (a) Partial (b) Positive (c) Neutral (d) Negative
- 37. Centistoke is the CGS unit for the following property
- (a) Surface tension
- (b) Coefficient of viscosity
- (c) Fluidity
- (d) Kinematic viscosity

- 38. Molar heat capacity of a system is defined as:
- (a) The amount of heat required to raise temperature of one mole of the system by one Kelvin
- (b) The amount of heat required to decrease temperature of one mole of the system by one Kelvin
- (c) The amount of heat required to decrease temperature of one mole of the system by one Fahrenheit
- (d) The amount of heat required to raise temperature of one mole of the system by one Fahrenheit
- 39. When a solid forms a gel more readily when gently shaken or otherwise sheared than when allowed to form the gel while the material is kept at rest, the phenomenon is known as
- (a) Thixotropy (b) Rheopexy (c) Negative rheopexy (d) Anti thixotropy
- 40. In the process of Extraction, ethanol is used as a solvent for
- (a) Sucrose
- (b) Waxes
- (c) Alkaloids
- (d) GUMS
- 41 Human Serum Albumin has a molecular weight of
- (a) 34,000
- (b) 65,000
- (c) 44,000
- (d) 59,000
- 42. Riboflavin, chemically is
- (a) 6,7-dimethyl isoalloxazine
- (b) 6,7-diethyl isoalloxazine
- (c) 8,9-dimethyl isoalloxazine
- (d) 8,9-diethyl isoalloxazine

- 43. Burow's solution is
- (a) Aluminium acetate solution strong
- (b) Calcium phosphate solution strong

(c) Ammonium acetate solution strong

(d) Calcium hydroxide solution

44. Drug products that contain the same therapeutic moiety but as different salts, esters or complexes are called as

(a) Therapeutic equivalents

(b) Pharmaceutical equivalents

(c) Pharmaceutical alternative

(d) Therapeutic alternatives

45. When six suppositories containing 20 percent of morphine hydrochloride in Theobroma oil are to be prepared, which of the following statement is correct? (Given: Displacement value of morphine hydrochloride is 1.5; weight of each suppository is 1 g)

- (a) The displacement value of morphine hydrochloride is to be considered
- (b) The displacement values of both morphine hydrochloride and Theobroma oil are to be considered
- (c) The displacement value of morphine hydrochloride is to be ignored
- (d) The displacement value of Theobroma oil is to be considered
- 46. Following amino acid does not exhibit optical isomerism:
- (a) Serine (b) Alanine (c) Glycine (d) Leucine

47. Fraction of administered dose (which may be -n form of an ester of salt) of active drug is termed as

(a) Activity factor (b) Formulation factor (c) Intrinsic factor (d) Salt factor

48. If two monosaccharides differ in their structure around a single carbon atom, they are called as

- (a) Epimers
- (b) Dimers
- (c) Trioses
- (d) Trimers

49. Which of the following is the example of "Invert Sugar"

- (a) Sucralose
- (b) Lactulose
- (c) Lactose
- (d) Sucrose

50. Which of the following groups stabilizes carbocation in electrophilic aromatic substitution

- (a) -CN, -SO3H
- (b) -COOH, -CHO
- (c) NH2, OH
- (d) -N(CH3)3+, -NO2

51. The mechanism of antiepileptic effect of Lamotrigine is by :

- (a) Inhibition of glutamate release
- (b) Blockade of NMDA receptors
- (c) Inhibition of Na+ channels
- (d) Inhibition of Ca++ channels

52. Closeness of a measured value to the true or accepted value is known as

(a) Accuracy (b) Deviation (c) Precision (d) Bias

53. Which of the following is Anti aromatic









54. Globar source in infra-red spectrophotometer is

- (a) Silicon carbide rod
- (b) Rhodium wire heated in ceramic
- (c) Tungsten filament
- (d) Nichrome wire

55. Cotrimoxazole is combination of trimethoprim with sulfamethoxazole. Which of the following statement is correct about Cotrimoxazole?
(a) Cotrimoxazole is contraindicated during pregnancy.
(b) Most effective ratio of trimethoprim: sulfamethoxazole in plasma is 20:1.
(c) Trimethoprim: sulfamethoxazole are administered in a dose ratio of 5:1 to achieve optimal synergistic effect.
(d) Cotrimoxazole is ineffective in treating respiratory tract infections.
56. Ion trapping is an important process in drug distribution with potential therapeutic benefit. It is defined as http://www.xamstudy.com
(a) An acidic drug will accumulate on the more basic side of the membrane and a basic drug on more basic side.
(b) An acidic drug will accumulate on the more acidic side of the membrane and a basic drug on more basic side.
(c) An acidic drug will accumulate on the more basic side of the membrane and a basic drug on more acidic side.
(d) An acidic drug will accumulate on the more acidic side of the membrane and a basic drug on more acidic side.
57. Identify the diuretic which causes hyperuricemia, tend to raise scrum calcium and also causes magnesium depletion
(a) Acetazolamide (b) Furosemide (c) Mannitol (d) Chlorthiazide
58. The Nitration of quinoline gives a mixture of
(a) 3-nitro-and 6-nitroquinolines (b) 2-nitro-and 7-nitroquinolines
(c) 5-nitro-and 8-nitroquinolines (d) 2-nitro-and 4-nitroquinolines
59. The specific reagents used m skraup synthesis of quinoline are
(a) Benzaldehyde, aniline, sulfuric acid and Nitrobenzene
(b) Glycerol, aniline, sulfuric acid and Nitrobenzene
(c) Aminoethanal and Nitrobenzene
(d) Glycerol, Acetal of aminoethanal and Nitrobenzene
60. Mitomycin C is an antibiotic isolated from
(a) Streptomyces peucetius (b) Streptomyces verticillus
(c) Streptomyces antibioticus (d) Streptomyces caespitosus
61. This 4-amino quinoline antimalarial drug marketed as R, S isomer, has two trifluromethyl moieties at position 2' and 8' and no electronegative substituent either at 6' or 7' positions is
(a) Mefloquine (b) Chloroquine (c) Primaquine (d) Amodiaquine
62. Which one is NOT an opioid receptor
(a) Gamma (b) Kappa (c) Mu (d) Delta
63. One of the following statement with respect to carbachol is NOT correct Identify:

- (a) It can be administered orally
- (b) It possess both muscarinic and nicotinic action
- (c) It is more susceptible to hydrolysis as compared to acetylcholine
- (d) The amino group decreases electrophilicity of the carbonyl group
- 64. What is the product of the following reaction?

$$\begin{array}{c|c}
 & NaCN, HCN \\
\hline
 & 80^{\circ}C-100^{\circ}C
\end{array}$$
(a)
$$\begin{array}{c|c}
 & NC \\
\hline
 & (c)
\end{array}$$
(b)
$$\begin{array}{c|c}
 & NC \\
\hline
 & (d)
\end{array}$$
(c)

- 65. Which three of the following are physiological effects of ganglionic stimulating agents
- (A) Stimulation of sympathetic ganglia and the adrenal medulla leading to cardiovascular responses
- (B) Initial transient stimulation and then a more persistent depression of all autonomic ganglia
- (C) Marked depression of CNS leading to analgesia.
- (D) Chronic exposure causes a marked increase in the density of nicotinic receptors contributing tolerance and dependence.
- (E) Combined activation of parasympathetic ganglia and cholinergic nerve ending by nicotine results in increased tone and motor activity of the bowel. (F) Inhibition of salivary and bronchial secretions that is followed by stimulation.
- (a) (A), (C) and (D)
- (b) (B), (D) and (F)
- (c) (B), (C) and (E)
- (d) (A); (B) and (D)
- 66. Two pore domain K+ ions channels are opened by which one of the following categories of anaesthetic
- (a) Barbiturates
- (b) Benzodiazepines (c) Opioid analgesics (d) Halogenated inhalation anaesthetic
- 67. Which of the following rearrangement involves migration of a group from carbon to electron deficient nitrogen
- (a) Wilgerodt rearrangement
- (b) Baeyer villiger, rearrangement
- (c) Pinacol-pinacolone rearrangement
- (d) Beckmann rearrangement
- 68. The mechanism by which fluorides inhibit dental caries is
- (a) By increasing susceptibility to acid
- (b) By increasing the sensitivity of tooth enamel
- (c) Decreased acid solubility of enamel
- (d) Increased acid solubility of enamel
- 69. The solvent used in NMR spectroscopy with the highest level of deuteriation among these is:

- (a) Deuterioacetone (b) Deuterum oxide (c) Deuteriomethanol (d) Deuteriochloroform

<ul> <li>70. The weakly acidic nature of warfarin is attributed to the presence of</li> <li>(a) Methylene protons on substitution at 3rd position</li> <li>(b) Proton of hydroxyl group at 4th position</li> <li>(c) Proton on chiral carbon</li> <li>(d) Aromatic protons of coumarin ring</li> </ul>					
71. Which of the following drugs possesses Antiplatelet effects (a) Erythropoietin (b) Urokinase (c) Desmopressin (d) Clopidogrel					
72. Walden inversion or complete reversal of stereochemistry occurs in one of the following substitution reaction (a) SNAr reactions (b) SNi reaction (c) SN2 reactions (d) SN1 reactions					
73. Antifungal agent 5-fluorocytosine is converted to its active metabolite 5-fluorouracil by one of the following fungal enzyme  (a) Cytosine deacetylase  (b) Cytosine reductase  (c) Cytosine demethylase  (d) Cytosine deaminase					
74. Propanone and 2-hydroxy propene are (a) Position isomers (b) Rotamers (c) Chain Isomers (d) Tautomers					
75. The principle of liquid-liquid extraction is (a) Partition (b) Solubility (c) Adsorption (d) Separation					
76. When an electron with charge $1.60 \times 10^{\circ}19$ coulombs is moved through a potential difference of 1 volt, the energy expended or released is called as					
77. An aldopentose can be converted into aldohexose by :					
(a) Ruffs degradation method (b) Wohls method					
(c) Killani-Fischer synthesis (d) Hoffmann degradation method					
78. Hydrated proton is called (a) Water proton (b) Hydronium ion (c) Roxonium ion (d) Proton pump					
79. What is the approximate chemical shift ranges for protons attached to aldehydic groups? (a) 4-8 $\delta$ (b) 9.5 - 10.0 $\delta$ (c) 6 - 7.9 $\delta$ (d) 1.5 - 3.5 $\delta$					
<ul> <li>80. The condensation of an active methylene compound with formaldehyde and an amine to form β-amino carbonyl compound is known as</li> <li>(a) Mannich reaction</li> <li>(b) Knoevenagel condensation</li> <li>(c) Stobbe condensation</li> <li>d) Beckmann rearrangement</li> </ul>					

(c) Preferential Cox-1 inhibitor		
(d) Selective Cox-2 inhibitor		
82. Which one of the following is not the p	roperty of osmotic diuretics	
(a) Undergo limited reabsorption by the rer	nal tubule	
(b) Are relatively inert pharmacologically		
(c) They inhibit carbonic anhydrase		
(d) Freely filtered at the glomerulus		
83. Electron releasing groups stabilizes free	e radicals by	
(a) Positive inductive ( + I) effect	(b) Negative mesomeric ( — m) effect	
(c) Positive mesomeric (+m) effect	(d) Negative inductive (-I) effect	
84. The principle of gas-liquid chromatogra	aphy is	
(a) Partition chromatography (b) Affinity chromatography		
(c) Adsorption chromatography	(d) Ion exchange chromatography	
85. The heterocyclic nucleus present in nitr	rofurantoin are	
(a) 3-nitro furan and imidazole 2, 4-dione		
(b) 2-nitro furan and imidazoline-2-one		
(c) Furan and pyrrole		
(d) 2-nitro furan and imidazole 2, 4-dione		
86. d-tubocurarine produces skeletal muscl	e relaxation by inhibiting	
(a) Ganglionic nicotinic receptors		
(b) Muscurinic receptors		
(c) Nicotinic receptors in neuromuscular ju	nction	
(d) alpha-adrenergic receptors		
	e triade effects of adrenaline leading to rise in blood pressure	
	creases the strength of ventricular contraction	
(b) An increased heart rate (positive chrono	•	
	specially in precapillary resistance vessels of skin	
(d) Stimulation of presynaptic alpha-2 adre Pharmacognosy	noreceptor leading to increase sympathetic tone Section :	
88. " Which one of the following is not the	biochemical marker of all death?	
(a) Phospholipase (b) Creatine kinase (c) L	ipase (d) Amylase	

89. In the transplant rejection the clinical features of graft-versus-host reaction includes following-Except

81. Meloxicam belongs to which class of Non-steroidal Anti-inflammatory Drugs (NSAIDs) (a) Preferential cox-2 inhibitor

(b) Selective Cox-1 inhibitor

(a) Dermatitis, diarrhoea

- (b) Fever, weight loss
- (c) Intestinal malabsorption
- (d) Rheumatoid arthritis
- 90. Which one of the following statement is true for cancer cells
- (a) Oncoproteins produced by cancer cells act a neighboring cells
- (b) Cancer cells require stimulation by growth factors (c) Cancer cells are highly sensitive to growth inhibitory signals
- (d) Cancer cells produce Oncoproteins in the absence of growth factors or external stimuli
- 91. Rosacea is a skin condition that affect:
- (a) Teenage people
- (b) Old age people
- (c) Only woman
- (d) Adults between the age of 30 60
- 92. A typical long bone consists of following parts except
- (a) The diaphysis (b) The trabeculae (c) The epiphyses (d) The periosteum
- 93. Which of the following Neurotransmitter is formed on demand and act immediately
- (a) Acetylcholine (b) Histamine (c) Nitric oxide (NO) (d) Adrenaline
- 94. In one of the following situations metabolic acidosis does not occur
- (a) Uncontrolled diabetes mellitus
- (b) Therapeutic administration of acetazolamide
- (c) Chronic renal failure
- (d) Production of less amount of lactic acid
- 95. Which one of the following is a organ specific (localized) autoimmune disease
- (a) Grave's disease (b) Scleroderma (c) Polyarteritis nodosa (d) Rheumatoid arthritis
- 96. One of the following is not a chemical or histochemical change that occurs in infarction of heart
- (a) Decrease in sodium
- (b) Glycogen depletion
- (c) Influx of calcium
- (d) Increase in lactic acid
- 97. Following pairs represent antibiotic and its source. Identify the correct pair
- (a) Griseofulvin: Bacteria
- (b) Polyenes: Fungi
- (c) Colistin: Fungi
- (d) Macrolides: Actinomycetes Section: Pharmacology
- 98. Who was the Chairman of the Drugs Enquiry Committee which was established in the year 1927?

- (a) B. Mukherjee (b) Joseph Bhore (c) S.S. Bhatia (d) R.N. Chopra
- 99. Which one of the following is not a renal osteodystrophy (type of bone disease associated with chronic kidney disease)?
- (a) Adynamic bone disease
- (b) Osteomalacia
- (c) Secondary hypoparathyroidism
- (d) Mixed renal osteodystrophy
- 100. Plants which are not differentiated into root, stem and leaves are :
- (a) Plant aginales (b) Principes (c) Thallophyte (d) Bromeliales
- 101. Identify antiarrhythmic drug that increases duration of QT wave and decreases the sinus rate, whereas it has no effect on QRS complex
- (a) Lidocaine (b) Amiodarone (c) Verapamil (d) Quinidine
- 102. As per the Drugs and Cosmetics Act-1940, if a drug is not labelled in prescribed manner it is a
- (a) Spurious drug (b) Substandard drug (c) Adulterated drug (d) Misbranded drug
- 103. "The major property of Ayurvedic herbs, "Rasa" indicates
- (a) Taste (b) Post digestive effect (c) Potency (d) Physicochemical properties
- 104. Except one of the following pairs represent drugs used in the treatment of glaucoma and their primary mechanism. Select the wrong pair from the following
- (a) Topical prostaglandin analogues: Increase aqueous outflow
- (b) Topical beta-adrenergic blockers: Decrease aqueous outflow
- (c) Topical miotics: Increase aqueous outflow
- (d) Topical carbonic anhydrase inhibitors: Decrease aqueous formation
- 105. Auto rhythmicity is present in which of the following group of tissues?
- (a) Skeletal muscle and cardiac muscle
- (b) Skeletal muscle and visceral smooth muscle
- (c) Cardiac muscle and neuronal tissue
- (d) Cardiac muscle and visceral smooth muscle
- 106. During the denaturation cycle in polymerase chain reaction the temperature is raised to:
- (a) 90 -98°C (b) 60 68°C (c) 70 78°C (d) 80 88°C
- 107. The general purpose sodalime glass is not a suitable material for fabricating the container for
- (a) Parenteral (b) Oral solutions (c) Liquids for external use (d) Dry powders
- 108. In which year India signed General Agreement on Trade and Tariffs (GATT) including Trade Related Intellectual Property Rights (TRIPS)
- (a) 1996 (b) 1994 (c) 1992 (d) 1990

- 109. A fixed oil when added to an equal volume of ethanol; clear liquid is obtained; on cooling at O°C and on storage for three hours, the liquid remains clear such fixed oil is identified as:
- (a) Castor Oil (b) Soyabean oil (c) Neem oil (d) Evening Primrose oil
- 110. In shikimic acid pathway, chorismate mutase converts chorismic acid to
- (a) Carotenoids (b) Phytol (c) Prephenate (Prephenic acid) (d) Gutta
- 111. Standards to be complied under D & C act -1940 for drugs imported, manufactured, stocked and exhibited for sale or distribution are covered under
- (a) Schedule M (b) Second Schedule (c) First Schedule (d) Schedule L
- 112. In colour test for alkaloids colchicine with mineral acids gives
- (a) Blue colour (b) Red colour (c) Yellow colour (d) Violet colour
- 113. Nicotine from tobacco is an alkaloid which is
- (a) Oxygen free liquid (b) Semisolid (c) Crystalline (d) Oxygen free solid
- 114. In solid-solid mixing, large scale continuous type of mixer is
- (a) Sigma blender
- (b) Ribbon blender
- (c) Zigzag blender
- (d) Twin shell blender
- 115. Which one of the following is not affected in respiratory tuberculosis
- (a) Pleural cavity (b) Mediastinal lymph nodes (c) Pericardium (d) Larynx
- 116. Identify phase of clinical trial having following features
- (A) Trial is conducted on about 3000 patients
- (B) Purpose of trial is therapeutic confirmation
- (C) Safety and tolerability is evaluated on wider scale
- (D) Completion of trial is followed by New Drug Application (NDA) Options –
- (a) Phase III (b) Phase I (c) Phase IV (d) Phase 11
- 117. Which of the following Microorganism is used for the effective synthesis of interferon
- (a) Micrococcus Luteus
- (b) Bacillus Subtilus
- (c) Saccharomyces Cerevisiae
- (d) Pseudomonas Aureginosa
- 118. In yam, the presence of, irregular arrangement of the fibres, the ends of which often project from the surface is because of
- (a) Absence of linters
- (b) Absence of combing
- (c) Presence of impurities

- (d) Improper drying
- 119. In International organization for standardization (ISO), standard pertaining to quality system model for quality assurance in production, installation and servicing is given under
- (a) ISO 9002 (b) ISO 9000 (c) ISO 9001 (d) ISO 9004
- 120. In Aloe the mucilage containing parenchymatous cells are present in
- (a) Central parenchymatous region
- (b) Pericyclic cells
- (c) Epidermis
- (d) Vascular bundles
- 121. Mechanical digestion in mouth in which food is mixed with saliva is and the movement of food from mouth into the stomach is
- (a) Mastication; deglutition
- (b) Deglutition; propulsion
- (c) Segmentation; mastication
- (d) Segmentation; propulsion
- 122. The Crystal form of Sulphacetamide is:
- (a) Rhombohedral (b) Orthorhombic (c) Monoclinic (d) Triclinic
- 123. As per the definition of D and C Act, Gudakhu (rubbed against human teeth) is considered as: (a) Food (b) Drug (c) Sweeting gum (d) Cosmetic
- 124. The fundamental principle "Law of similia" falls under which therapy
- (a) Ayurveda (b) Siddha
- (c) Homoeopathy
- (d) Aroma therapy
- 125. Dragendorff's reagent is
- (a) Potassium iodate
- (b) Potassium bismuth iodide
- (c) Potassium mercuric iodide
- (d) Potassium picrate

**GPAT 2020 ANSWERS KEY** 

	(4					•		
1 - b	16 - d	31 - a	46 - с	61 - a	76 - d	91 - d	106 - a	121 - a
2 - c	17 - d	32 - a	47 - d	62 - a	77 - c	92 - b	107 - a	122 - b
3 - d	18 - a	33 - b	48 - a	63 - c	78 - b	93 - с	108 - b	123 - d
4 - c	19 - b	34 - a	49 - d	64 - a	79 - b	94 - d	109 - a	124 - с
5 - b	20 - a	35 - b	50 - c	65 - d	80 - a	95 - a	110 - с	125 - b
6 - c	21 - b	36 - d	51 - c	66 - d	81 - a	96 - a	111 - b	
7 - c	22 - c	37 - d	52 - a	67 - d	82 - c	97 - d	112 - с	
8 - c	23 - d	38 - a	53 - d	68 - c	83 - a	98 - d	113 - a	
9 - c	24 - d	39 - b	54 - a	69 - a	84 - a	99 - c	114 - с	
10 - c	25 - c	40 - c	55 - a	70 - b	85 - d	100 - c	115 - с	
11 - b	26 - b	41 - b	56 - с	71 - d	86 - c	101 - b	116 - a	
12 - a	27 - с	42 - a	57 - d	72 - c	87 - d	102 - d	117 - с	
13 - a	28 - a	43 - a	58 - с	73 - d	88 - a	103 - a	118 - b	
14 - b	29 -d	44 - c	59 - b	74 - d	89 - d	104 - b	119 - a	
15 - b	30 - d	45 - c	60 - d	75 - a	90 - d	105 - d	120 - a	

Everything is a remix